Draft version January 20, 2013
Preprint typeset using LaTeX style emulateapj v. 03/07/07

arXiv:1105.0883v1 [astro-ph.HE] 4 May 2011

AN ANOMALOUS QUIESCENT STELLAR MASS BLACK HOLE

Mark T. Reynolds[1], Jon M. Miller[1]

Draft version January 20, 2013

ABSTRACT

We present the results of a 40 ks *Chandra* observation of the quiescent stellar mass black hole GS 1354-64. A total of 266 net counts are detected at the position of this system. The resulting spectrum is found to be consistent with the spectra of previously observed quiescent black holes, i.e., a power-law with a photon index of $\Gamma \sim 2$. The inferred luminosity in the $0.5-10$ keV band is found to lie in the range $0.5-6.5 \times 10^{34}$ erg s^{-1}, where the uncertainty in the distance is the dominant source of this large luminosity range. Nonetheless, this luminosity is over an order of magnitude greater than that expected from the known distribution of quiescent stellar mass black hole luminosities and makes GS 1354-64 the only known stellar mass black hole to disagree with this relation. This observation suggests the possibility of significant accretion persisting in the quiescent state.

Subject headings: accretion, accretion discs - black hole physics - stars: binaries (GS 1354-64) X-rays: binaries

1. INTRODUCTION

Observations at optical and infrared wavelengths have allowed us to measure the mass of 19 Galactic and 4 extra-galactic stellar mass black holes ($M_{BH} \sim 10-20 M\odot$: McClintock & Remillard 2006; Orosz et al. 2007; Silverman et al. 2008; Crowther et al. 2010; Corral-Santana et al. 2011). The presence of a black hole is inferred whenever the mass of the compact object is dynamically constrained to be greater than 3 $M_\odot$ (Rhoades & Ruffini 1974; Kalogera & Baym 1996). This does *not* confirm the compact object to be a classical black hole and these observations could be interpreted to agree with more exotic physics (see Narayan & McClintock 2008 for details). In order to confirm that the objects above are indeed black holes, one is required to detect a unique characteristic of such objects, i.e. the event horizon.

X-ray observations of accreting black holes have revealed a number of distinct accretion regimes, which correlate in a broad sense with luminosity of the source. The very-high, soft and hard states (hereafter VHS, SS & HS) are the primary active accretion states observed in XRBs. At luminosities below that observed in the HS state the system is said to be in the quiescent state. The quiescent state is characterized as an extremely faint state ($L_x \leq 10^{33.5}$ erg s^{-1}), with an X-ray spectrum that is distinctly non-thermal ($\Gamma = 1.5 - 2.2$; McClintock & Remillard 2006; Corbel et al. 2006).

The quiescent X-ray emission from black hole binaries (BHBs) is not consistent with expectations from standard accretion disc theory (e.g. McClintock et al. 1995). The current paradigm for understanding the X-ray emission from quiescent BHBs involves a standard thin disc which transforms to a quasi-spherical inner flow at a distance of $\sim 10^3 - 10^4$ Schwarzschild radii from the black hole. The inner flow could consist of an advection dominated accretion flow (ADAF: see Narayan & McClintock 2008 and references therein). In an ADAF the energy, released via viscous dissipation, remains in the accreting gas rather than being radiated away. As a result, most of the energy is advected with the accretion flow resulting in only a small percentage of the energy being radiated by the gas before it reaches the compact object. The radiative efficiency of such a flow is expected to be in the range $\sim 0.01 - 1\%$. In comparison a standard thin-disc is expected to have an efficiency an order of magnitude greater. The ADAF model has been successfully applied to observations of a number of quiescent and hard state BHBs, e.g. Narayan et al. (1996). The ADAF solution is not limited to stellar mass black holes and has also been successfully used to model the emission from a number of supermassive black holes (SMBH), e.g. see Yuan et al. (2003) for an ADAF fit to the spectrum of the least luminous known black hole Sgr A* ($L_x \sim 10^{-9}$ L_{Edd}).

However, the ADAF solution also allows outflows as emphasized by Blandford & Begelman (1999). In recent years jets/outflows have been recognized as a ubiquitous feature associated with the process of accretion on the largest and smallest scales, e.g. active galactic nuclei, XRBs. Fender et al. (2003) proposed an alternative scheme, whereby at low luminosity ($L_x \lesssim 10^{-4}$ L_{Edd}) BHs should enter a 'jet-dominated' state, in which the majority of the accretion power drives a radiatively-inefficient jet. The detection of the black hole A0620-00 at radio wavelengths may support this (Gallo et al. 2006), although the absence of a discernible jet in Sgr A* is problematic (Narayan & McClintock 2008). However, the detection of frequency dependent time-lags in the radio flares from Sgr A* (Yusef-Zadeh 2009), combined with the stratified nature of the radio and mm-wave emission (Doeleman et al. 2008) strongly suggest an unbound, mildly relativistic outflow. Recent work envisages the X-ray flux from a quiescent system to be a combination of an outflow/jet and an inner advective region, e.g. Yuan et al. (2009).

With the advent of the *Chandra* & *XMM-Newton* X-ray observatories detailed observations of quiescent black hole and neutron star binary systems have become possible, i.e. $L_x \lesssim 10^{-6}$ L_{Edd}. As first pointed out in Narayan et al. (1997) & Garcia et al. (1998), the ob-

---

Electronic address: markrey@umich.edu
[1] Department of Astronomy, University of Michigan, 500 Church Street, Ann Arbor, MI 48109



FIG. 1.— GS 1354-64 background subtracted lightcurve as observed by *Chandra*, where the lightcurve has been divided into 5 ks bins. Variability of a factor of two is observed on kilo-second timescales.

served luminosities of the quiescent BHs are systematically fainter than NSs. For a black hole the energy stored in the flow is advected across the event horizon whereas for a neutron star the material strikes the solid surface where it is re-radiated (Garcia et al. 2001; Kong et al. 2002). Subsequent observations have confirmed this picture (Sutaria et al. 2002; Hameury et al. 2003; Tomsick et al. 2003; McClintock et al. 2004; Corbel et al. 2006; Gallo et al. 2008).

GS 1354-64 was discovered by *Ginga* in 1987, where it displayed an X-ray spectrum dominated by a soft blackbody disc component, consistent with that of the then known black hole binaries in the soft state (Makino 1987; Kitamoto et al. 1990). A second outburst was observed from this system in 1997 by *RXTE*, though on this occasion the system remained in the low-hard state throughout the outburst (Brocksopp et al. 2001). The black hole nature of the primary has been dynamically confirmed by Casares et al. (2004, 2009), who found the system to comprise of a G0-5 III mass donor in a $\sim$ 2.5 days orbit around the black hole. The measured radial velocity ($\sim$ 279 km s^{-1}) in combination with the absence of any X-ray eclipses sets a secure lower limit for the mass of the black hole of $M_x \geq 7.6 \pm 0.7\ M_\odot$. GS 1354-64 is found to lie at a large distance, where 25 kpc $\leq$ d $\leq$ 61 kpc, and the upper limit is obtained by requiring the 1987 outburst to be sub-Eddington. This makes GS 1354-64 the most distant known Galactic black hole. The field containing GS 1354-64 has not been previously observed by either *XMM-Newton* or *Chandra*. The only X-ray imaging of this field was carried out by *Rosat* as part of the all sky survey (Voges et al. 1999), where only a weak upper limit exists, $f_x \leq 5.6 \times 10^{-13}$ erg s^{-1} cm^{-2}.

In this letter, we describe an observation of the quiescent Galactic stellar mass black hole binary GS 1354-64 with the *Chandra X-ray Observatory*. We describe the observations and extraction of source spectra and lightcurves, these data are discussed in the context of models for the quiescent accretion flow.

2. OBSERVATIONS

GS 1354-64 was observed by *Chandra* for 40 ks on 2010 October 1 (MJD 55470, obsid: 12471, PI: Reynolds), where it was placed on the back illuminated ACIS-S3 detector, which was operated in VFAINT mode. A significant point source is detected consistent with the known position of GS 1354-64 (Brocksopp et al. 2001). As the I3, S1–S4 detectors were active during this observation, a number of serendipitous point sources are also detected. These sources (31 in total) and their properties will be described in a separate publication.

The GS 1354-64 spectrum was extracted from a 2.5″ region with `psextract`, which also generated the appropriate response files. A background spectrum was extracted from a neighbouring source free region on the detector. We detect 266 net counts consistent with the known position of GS 1354-64. The source & background lightcurves were extracted using `dmextract` and binned using `lcurve`. Inspection of the background lightcurve reveals an absence of any significant flares during the observation. The background subtracted lightcurve is displayed in Fig. 1, where it has been divided into 5 ks bins. Variability of a factor of 2x is observed on kilo-second timescales, e.g., the count rate is observed to increase from an average of $\sim$ 0.006 ct s^{-1} to $\sim$ 0.012 ct s^{-1} at the 30 ks mark.

All data reduction and analysis takes place within the HEASOFT 6.6.2 environment, which includes FTOOLS 6.6, CIAO 4.2 and XSPEC 12.5.0AJ. The latest versions of the relevant *Chandra* CALDB files are also used.

3. ANALYSIS

The relatively large number of counts detected facilitates basic spectral fitting. The spectrum was binned such that each bin contained 20 counts using `grppha`, providing useful data in the spectral range 1 – 6 keV. We model the spectrum using 2 models for the continuum (i) a power-law (`pha*po`), and (ii) Bremsstrahlung (`pha*brem`). The results of the model fits are displayed in Table. 1. The best fit power-law spectral index of $\Gamma \sim 2.1$ (see Fig 2) and the bremsstrahlung temperature of $T_B \sim 5$ keV are consistent with previous observations of quiescent stellar mass black holes, e.g., Kong et al. (2002); Bradley et al. (2007). We measure an absorbed flux of $\sim$ 7.7(7.4) $\times$ 10^{-14} erg s^{-1} cm^{-2} in the 0.5 – 10.0 keV band for the power-law(bremsstrahlung) model. This corresponds to an unabsorbed flux of approximately $\sim 1.5(1.1)\times10^{-13}$ erg s^{-1} cm^{-2} in the same energy range, see Table 1 for details.

In Fig. 3, we plot contours of power-law index and bremsstrahlung temperature versus extinction for each of the models above. The uncertainty regions are large consistent with the small energy range covered by the data. Nonetheless, we see that the power-law model favours significantly larger values for the extinction $N_H \sim 9 \times 10^{21}$ cm^{-2}. For comparison, Kitamoto et al. (1990) estimated E(B-V) $\sim$ 1 ($N_H \sim 5.3 \times 10^{21}$ cm^{-2}), and the extinction estimated from the optical spectra obtained by Casares et al. (2004, 2009) are consistent with this value.

As the measured spectrum is intrinsically soft, the differing extinction estimates result in the unabsorbed flux for the power-law model being approximately 1.5 times that measured using the bremsstrahlung model. The power-law model is formally a better fit to the observed spectrum; however, the low number of bins render this insignificant.



FIG. 2.— *Chandra* 1 – 6 keV spectrum of GS 1354-64. The exposure time is ~ 39 ks. The best fit power-law model ($\Gamma \sim 2.1$) is plotted, the source luminosity is $L_x \sim 0.7 - 6 \times 10^{34}$ erg s^{-1}, see Table 1 for details.

In Fig. 4, we plot the luminosity calculated using the unabsorbed flux from the best fit power-law model in Table 1 and the distance estimates of Casares et al. (2009), i.e., $25 \leq d_{kpc} \leq 61$. GS 1354-64 is observed to be at a luminosity over an order of magnitude greater than that expected from a quiescent black hole. For this system to be consistent with the known distribution of black holes would require a decrease of the distance to ~ 5.8 kpc. Monitoring observations at optical wavelengths with the Faulkes telescope[2] revealed GS 1354-64 to be consistent with the known quiescent optical flux at the time of our *Chandra* observations, though, we note the nearest observation took place approximately 2 months prior to the *Chandra* observation described herein. While observations with the available X-ray all sky monitors show no evidence for increased activity either before or after our observation. However, a faint outburst would not have been detected by these observations.

There are 2 other stellar mass black holes with weak quiescent luminosity constraints. The first system, H 1705-250, had only been observed by *Rosat*. This upper limit will be updated soon by *Chandra* (Obsid: 11041, PI: Kong). The second system, GX 339-4, is an extremely active transient having exhibited 5 large and ~ 10 small outburst in the last 15 years. The most sensitive observation was obtained by *Chandra* in 2003 and revealed this system at the faintest level measured to date. However, this observation occurred ~ 10 months after the 2002 outburst and 7 months prior to the 2003/2004 outburst (Gallo et al. 2003). Hence, it is highly unlikely that GX 339-4 was in quiescence at the time of this observation. The current upper limits for these systems are represented by the solid triangles in Fig. 4.

4. DISCUSSION

We have observed the quiescent stellar mass black hole GS 1354-64 with *Chandra* and found it to be at a luminosity of ~ 10^{34} erg s^{-1}. This is over an order of magnitude greater than expected. Here, we discuss the uncertainty in the luminosity estimate and consider the



FIG. 3.— Contour plots corresponding to the best fit models in Table 1. The power-law fit is represented by the solid lines, while the thermal bremsstrahlung model is represented by the dotted contours. Contours plotted represent the 68% and 90% confidence levels.

implications of this measurement for the nature of the accretion flow in the quiescent state.

4.1. *Distance to GS 1354-64*

In order to compare to the other systems in Fig. 4, we must know the distance to each system. Casares et al. (2004, 2009) have calculated the minimum distance to GS 1354-64 to be $d_{min} \geq 25$ kpc. The distance estimate depends on the known spectral type of the secondary star, the apparent magnitude of the star, the contribution from the accretion disk and the extinction. The extinction is the most uncertain quantity in the distance calculation. Casares et al. (2004) estimate the extinction from known empirical relationships between observed spectral features and the column density resulting in an estimated E(B-V) ~ 1, or more specifically E(B-V) ≥ 0.78, consistent with the estimate of Kitamoto et al. (1990). However, the measured neutral hydrogen column in this field suggests a larger value, i.e., $N_H \sim 7.27 \times 10^{21}$ cm^{-2} (Kalberla et al. 2005) or E(B-V) ~ 1.4.

A higher level of extinction will effect the estimated minimum distance to this system of $d_{min} \geq 25$ kpc, in particular for a column density of $N_H \sim 7.27 \times 10^{21}$ cm^{-2} we find a minimum distance of $d_{min} \geq 17.8$ kpc, decreasing to 10.2 kpc for a column of 10^{22} cm^{-2}. The empirical diffuse interstellar band relation of Herbig (1975) as used in Casares et al. (2004) suggests an upper limit to the reddening of E(B-V) ≤ 1.5 ($N_H \sim 7.9 \times 10^{21}$ cm^{-2}) corresponding to a distance of ~ 15.6 kpc. Nonetheless, even in the unlikely event that the extinction is underestimated, GS 1354-64 remains too luminous in comparison to the known distribution.

4.2. *Possibilities*

The measured X-ray luminosity allows one to estimate the mass accretion rate, we do this under 2 assumptions for the accretion efficiency, i.e., $L_x = \eta \dot{M} c^2$. Assuming standard thin disc accretion ($\eta = 0.1$) implies an accretion rate of 10^{-11} ($L_x/10^{34}$) $M_\odot$ yr^{-1}, whereas for an ADAF an efficiency of $\eta = 10^{-2} - 10^{-4}$ is expected, and hence, a commensurably larger accretion rate. Accretion rates onto the outer disc of ~ 10^{-10} $M_\odot$ yr^{-1} have been estimated for a number of quiescent black hole binaries,

---

[2] `http://staff.science.uva.nl/~davidr/faulkes/` (Lewis et al. 2008)



Fig. 4.— The vertical blue line indicates the measured luminosity of GS 1354-64 assuming a distance of 25 kpc ≤ d ≤ 61 kpc (Casares et al. 2009), while the dotted extension indicates the estimated lower limit to the distance, see §4.1 for details. GS 1354-64 is observed to be over an order of magnitude too luminous in comparison to the known distribution of quiescent black holes (solid blue symbols). The open red symbols indicate the quiescent luminosities of neutron stars. Upper limits are represented by triangles. The red and blue filled areas indicate the regions in the L_x, P_{orb} plane occupied by the neutron star and black holes systems respectively. Black hole luminosities are from Garcia et al. (2001); Kong et al. (2002); Sutaria et al. (2002); Tomsick et al. (2003); Hameury et al. (2003); McClintock et al. (2004); Corbel et al. (2006); Gallo et al. (2008), and the neutron star luminosities are from Garcia et al. (2001); Tomsick et al. (2004); Lin et al. (2009); Degenaar et al. (2011).

e.g., A0620-00 (McClintock et al. 1995) and XTE J1550-564 (Orosz et al. 2011). As such, the accretion rates necessary to generate the observed luminosity appear to be feasible.

We also note the similarity between the observed X-ray variability of a factor of ∼ 2 (see Fig. 1) and that typically observed at optical wavelengths, where variability of approximately 1 magnitude (∼ 2.5x in linear units) is common from GS 1354-64 in quiescence (Casares et al. 2004, 2009). Correlated optical/X-ray variability has been observed from V404 Cyg in quiescence. However, the origin of this variability is not understood, with emission from the inner ADAF and/or irradiation of the disk being likely (Hynes et al. 2004). A similar scenario is likely here, in the case of GS 1354-64; however, further observations are required to confirm this possibility.

Below we briefly discuss what we consider to be the most likely explanation for the large observed quiescent luminosity:

(i) A temporary increase in the mass accretion rate, $\dot{M}$. For example, observations of the stellar mass black hole GRO J1655-40 ($P_{orb} \sim 63$ hrs) have revealed luminosity variations by a factor of 10 (Hameury et al. 2003). However, the brighter of these observations was obtained ∼ 9 months after the 1995 outburst and only one month prior to the 1996 outburst, i.e., the observed large luminosity was caused by the system not being quiescent at the time of the *ASCA* observations. Long term variations of a factor of ∼ 10 have also been observed from V404 Cyg ($P_{orb} \sim 155$ hrs, Bradley et al. 2007), while variations of ∼ 20x have been observed on kilo-second timescales with *Chandra* (Hynes et al. 2004). However, in both cases the luminosity remained consistent with that expected for the distribution of quiescent black holes, see Fig. 4. Observations of Sgr A* have revealed flares with an amplitude of greater than 100 times the true quiescent rate (Baganoff et al. 2001; Porquet et al. 2008), but these flares typically have a duration of a few hours and as such are unlikely the cause of the large luminosity, which we observe from GS 1354-64.

(ii) The actual quiescent accretion luminosity from this system is $\gtrsim 10^{34}$ erg s^{-1}. This would be the largest Eddington scaled luminosity measured from a black hole in quiescence to date, and would point to the existence of a unique low luminosity accretion flow in this system. Standard accretion disc theory predicts the disk to be truncated, with a low mass accretion rate and luminosity (McClintock et al. 1995; Narayan et al. 1996; Lasota et al. 2008), e.g., both GRO J1655-40 ($M_x \sim 6.5\ M_\odot$, $P_{orb} \sim 63$ hrs) & V4641 Sgr ($M_x \sim 7\ M_\odot, P_{orb} \sim 68$ hrs) have similar orbital periods to GS 1354-64 ($M_x \gtrsim 7\ M_\odot, P_{orb} \sim 60$ hrs), but have quiescent luminosities consistent with the other black holes, i.e., $L_x \sim 5.9 \times 10^{31}$ erg s^{-1}, 4×10^{31} erg s^{-1} respectively (Hameury et al. 2003; Tomsick et al. 2003). The nature of the mass donor secondary star is also sufficiently similar with ∼ F4III-IV (Orosz & Bailyn 1997) and ∼ B9III (Orosz et al. 2001) respectively in comparison to the G0-G5III secondary in GS 1354-64. If the large quiescent luminosity we have measured is confirmed, it would suggest the existence of a previously unrecognized stable mode of low luminosity accretion ($L_x \sim 10^{-5}\ L_{Edd}$), and with it a population of relatively faint accreting black holes in the galaxy, e.g., Menou et al. (1999).

4.3. *BH vs NS luminosities*

A comparison of the quiescent luminosities of the black hole and neutron star binaries revealed the black hole systems to be a factor of ∼ 100 times fainter (Garcia et al. 2001, Fig. 4). This has been interpreted as evidence for the absence of a solid surface in the black hole systems, and conversely as indirect evidence for the existence of the event horizon. The observation presented of GS 1354-64 herein contradicts this empirical relationship. Previously, an observation of the neutron star binary 1H 1905+00 was claimed to also contradict this relationship (Jonker et al. 2007); however, further analysis revealed this to be false (Lasota et al. 2008). If the measured luminosity of GS 1354-64 (Fig. 4) is shown to be stable by future observations, one could ask if the observed difference between the black hole and neutron stars in quiescent is not caused by the absence of a surface in the black hole systems but instead by the favoring of this higher luminosity quiescent accretion flow in the neutron star systems.

We thank Dave Russell for discussions regarding the Faulkes monitoring of this system. MTR & JMM gratefully acknowledge support through the *Chandra* Guest Observer program. This research made use of the *SIMBAD* database, operated at CDS, Strasbourg, France and NASA's Astrophysics Data System. We thank the anonymous referee for useful comments

TABLE 1
BROADBAND CONTINUUM FIT PARAMETERS

Model	N_H [10^{22} cm^{-2}]	Γ	kT [keV]	$f_{x\ abs}(0.5 - 10.0$ keV) [10^{-14} erg s^{-1} cm-2]	$f_x(0.5 - 10.0$ keV) [10^{-14} erg s^{-1} cm-2]	L_x(25kpc, 61kpc) [10^{33} erg s^{-1}]	χ^2/ν
`pha(po)`	$0.9^{+0.45}_{-0.4}$	$2.2^{+0.6}_{-0.4}$	–	$7.69^{+0.9}_{-0.7}$	$14.5^{+1.7}_{-1.3}$	$7.65^{+0.90}_{-0.68}$, $64.6^{+7.5}_{-5.8}$	14.3/10
`pha(bremss)`	$0.58^{+0.34}_{-0.28}$	–	$5.1^{+7.3}_{-2.4}$	$7.4^{+0.9}_{-0.7}$	$10.7^{+1.0}_{-1.2}$	$5.65^{+0.52}_{-0.64}$, $47.6^{+4.5}_{-5.3}$	15.7/10

NOTE. — Best fit model parameters for the GS 1354-64 continuum as measured on the ACIS-S3 detector in the spectral range 1 – 6 keV, see Fig. 2. The flux is calculated in the 0.5 – 10.0 keV band, and the luminosity has been calculated assuming a distance of $25 \leq d_{kpc} \leq 61$ (Casares et al. 2009). All errors are quoted at the 90% confidence level.

REFERENCES

Baganoff F.K., Batz M.W., Brandt W.N. et al., 2001, Natur, 413, 45

Blandford R., Begelman M., 1999, MNRAS, 303, 1

Bradley C.K., Hynes R.I., Kong A.K.H., Haswell C.A., Casares J., Gallo E., 2007, ApJ, 667, 427

Brocksopp C., Jonker P.G., Fender R.P., Groot P.J., van der Klis M., Tingay S.J., 2001, MNRAS, 323, 517

Casares J., Zurita C., Shahbaz T., Charles P.A., Fender R.P., 2004, ApJ, 613, 133

Casares J., Orosz J.A., Zurita C. et al., 2009, ApJS, 181, 238

Corbel S., Tomsick J.A., Kaaret P., 2006, ApJ, 636, 971

Corral-Santana J.M., Casares J., Shahbaz T., Zurita C., Martnez-Pais I.G., Rodrguez-Gil P., 2011, MNRAS, 413, 15

Crowther P.A., Barnard R., Carpano S., Clark J.S., Dhillon V.S., Pollock A.M.T., 2010, MNRAS, 403, 41

Degenaar N., Wolff M.T., Ray P.S. et al., 2011, MNRAS, 412, 1409

Doeleman S.S., Weintroub J., Rogers A.E.E. et al., 2008, Natur, 455, 78

Fender R.P., Gallo E., Jonker P.G., 2003, MNRAS, 343, 99

Gallo E., Fender R., Corbel S., 2003, ATel, 196

Gallo E., Fender R.P., Miller-Jones J.C.A., Merloni A., Jonker P.G., Heinz S., Maccarone T., Van der Klis M., 2006, MNRAS, 370, 1351

Gallo E., Homan J., Jonker P.G., Tomsick J.A., 2008, ApJ, 683, 51

Garcia M.R., McClintock J.E., Narayan R., Callanan P.J., 1998, ASPC, 137, 506

Garcia M.R., McClintock J.E., Narayan R., Callanan P., Barret D., Murray S., 2001, ApJ, 553, 47

Hameury J.M., Barret D., Lasota J.P., McClintock J.E., Menou K., Motch C., Olive J.F., Webb N., 2003, A&A, 399, 631

Herbig G.H., 1975, ApJ, 196, 129

Hynes R.I., Charles P.A., Garcia M.R. et al., , 2004, ApJ, 611, 125

Jonker P.G., Steeghs D., Chakrabarty D., Juett A.M., 2007, ApJ, 665, 147

Kalberla P.M.W., Burton W.B., Hartmann D., Arnal E.M., Bajaja E., Morras R., Pöppel W.G.L., 2005, A&A, 440, 775

Kalogera V., Baym G., 1996, ApJ, 470, 61

Kitamoto S., Tsunemi H., Pedersen H., Ilovaisky S.A., van der Klis M., 1990, ApJ, 361, 590

Kong A.K.H., McClintock J.E., Garcia M.R., Murray S.S., Barret D., 2002, ApJ, 570, 277

Lasota J.P., 2008, NewAR, 51, 752

Lewis F., Roche P., Russell D.M., Fender R.P., 2008, AIPC, 1010, 204

Lin J., Nowak M.A., Chakrabarty D., 2009, ApJ, 706, 1069

Makino, 1987, IAUC, #4342

McClintock J.E., Horne K., Remillard R.A., 1995, ApJ, 442, 358

McClintock J.E., Narayan R., Rybicki G.B., 2004, ApJ, 615, 402

McClintock J.E., Remillard R.A., 2006, in Lewin W.H.G., van der Klis M., eds, Compact Stellar X-Ray Sources, Cambridge University Press, Cambridge

Menou K., Narayan R., Lasota J.P., 1999, ApJ, 513, 811

Narayan R., McClintock J.E., Yi I., 1996, ApJ, 457, 821

Narayan R., Garcia M.R., McClintock J.E., 1997, ApJ, 478, 79

Narayan R., McClintock J.E., 2008, NewAR, 51, 733

Orosz J.A., Bailyn C.D. 1997, ApJ, 477, 876

Orosz J.A., Kuulkers E., van der Klis M. et al., 2001, ApJ, 555, 489

Orosz J.A., McClintock J.E., Narayan R., 2007, Natur, 449, 872

Orosz J.A., Steiner J.F., McClintock J.E., Torres M.A.P., Remillard R.A., Bailyn C.D., Miller J.M., 2011, ApJ, 730, 75

Porquet D., Grosso N., Predehl P. et al., 2008, A&A, 488, 549

Rhoades C.E., Ruffini R., 1974, PhRvL, 32, 324

Silverman J.M., Filippenko A.V., 2008, ApJ, 678, 17

Sutaria F.K., Kolb U., Charles P. et al., 2002, A&A, 391, 993

Tomsick J.A., Corbel S., Fender, R. et al., 2003, ApJ, 597, 133

Tomsick J.A., Gelino D.M., Halpern J.P., Kaaret P., 2004, ApJ, 610, 933

Voges W., Aschenbach B., Boller Th. et al., 1999, A&A, 349, 389

Yuan F., Quataert E., Narayan R., 2003, ApJ, 598, 301

Yuan F., Yu Z., Ho L.C., 2009, ApJ, 703, 1034

Yusef-Zadeh F., Bushouse H., Wardle M. et al., 2009, ApJ, 704, 348

This paper was typeset using a LaTeX file prepared by the author